arXiv:0810.2514v1 [math.AP] 14 Oct 2008

UNIQUENESS OF SELF-SIMILAR SOLUTIONS TO THE NETWORK FLOW IN A GIVEN TOPOLOGICAL CLASS

MARIEL SÁEZ TRUMPER

Abstract. In this paper we study the uniqueness of expanding self-similar solutions to the network flow in a fixed topological class. We prove the result via the parabolic Allen-Cahn approximation proved in [S1].

Moreover, we prove that any regular evolution of connected tree-like network (with an initial condition that might be not regular) is unique in a given a topological class.

1. Introduction

The flow of planar networks by curve shortening flow has been of interest for several authors in last few years (see [MNT], [Sch], [SS] and [MS] for example). A planar network in $\Omega \subset \mathbb{R}^2$ is a finite union of arcs embedded on the plane such that each pair of curves may intersect each other only at their ends. Moreover, these ends always intersect either other arc or $\partial\Omega$. These intersections are called the vertices of the network.

A network is said to evolve by curve shortening flow if the evolution set is a network for every fixed time and each of its constituent arcs γ_i satisfies the shortening flow equation:

$$\frac{d\gamma_i}{dt} \cdot \nu = k_{\gamma_i}.$$

In order to have a well defined equation it is necessary to impose other conditions at the interior nodes. The most natural condition in the context of Brakke's work ([Br]) is to impose that the interior vertices are all trivalent and that the meeting angles of the curves are all equal to $\frac{2\pi}{3}$. When a network satisfies this condition during its evolution we say that the network is regular. We would like to remark that it is possible to impose other conditions for the angles at the interior nodes, but these will not be consider in this paper. At the exterior nodes we will impose Dirichlet boundary condition, namely we will prescribe the evolution of these nodes (mostly considering them to be fixed).

Recently, in [MS] was proved that for an initial condition of k half-lines there exists a connected solution to the network flow. Moreover, this solution is regular for all positive times and self-similar; however, it is not necessarily unique. In this paper we discuss uniqueness for these connected networks within a topological class.

To define the topological class we assume that the exterior nodes are fixed. If such vertices do not exist, that is if the curves extend to infinity, by "fixing the vertices" we mean that at infinity the network is asymptotic to certain fixed curves (which, in the case of the self-similar solutions described above, agree with the k half-lines that were taken as initial condition).

Definition 1.1. *We say that two networks belong to the same topological class if there is a homotopy between them relative to the exterior vertices.*

The author was supported by Proyecto Fondecyt de Iniciación 11070025.

As an example of this concept in Figure 1(a) we show two networks in the same topological class, which contrast with Figure 1(b), where the two networks are in different topological classes. We remark that although in certain situations the networks in Figure 1(b) can be in the same topological class after rotation, we do not allow this by fixing the ends in the homotopy process.



(a) Two networks of the same topological class (b) Two networks of the different topological class

Figure 1. Topological Class of networks with 4 exterior nodes

Now we can state the main theorem of this paper:

Theorem 1.1. *For each topological class there is at most one connected self-similar solution to the network flow with initial condition consisting of k lines meeting at the origin and that is regular for positive times.*

This theorem can be extended as follows:

Theorem 1.2. *Suppose that we have an arbitrary connected tree-like network that has an evolution via network flow in a domain Ω that is regular for $0 < t < T$ and that at the boundary satisfy Dirichlet conditions. Moreover, assume that the curvatures k^i of this network are bounded by $\frac{C}{\sqrt{t}}$. Then, given a topological class, there is a unique evolution for this network.*

We would like to remark that for unbounded domains Ω we assume that the boundary conditions satisfied by the networks above are analogous to the ones satisfied by the self-similar solutions. Namely, we assume that the arcs converge strongly (at least in $C^{2,1}$) to a fixed curve, which is assumed to be compatible with the evolution equation (that in the case of the self-similar solutions described earlier, correspond to lines and in general could be any unbounded curve that is a solution to the curve shortening flow equation).

The evolution of networks can be also seen from a different point of view, namely as the nodal set of the limit of solutions to the *vector-valued parabolic Allen-Cahn* equation. The

Allen-Cahn Equation (with Dirichlet boundary condition) is given in a domain Ω by:

$$\frac{\partial u_\epsilon}{\partial t} - \Delta u_\epsilon + \frac{\nabla_u W(u_\epsilon)}{\epsilon^2} = 0 \text{ for } x \in \Omega \tag{1}$$

$$u_\epsilon(x,0) = \psi_\epsilon(x), \tag{2}$$

$$u_\epsilon|_{\partial\Omega} = \phi_\epsilon(x,t) \tag{3}$$

where $u_\epsilon : \mathbb{R}^n \times \mathbb{R}_+ \to \mathbb{R}^m$ and $W : \mathbb{R}^n \to \mathbb{R}$ is a positive potential with a finite number of minima. In particular we will concentrate on the case $m = n = 2$ and W is a function with 3 minima.

In [BR] Bronsard and Reitich studied formally this equation and predicted that as $\epsilon \to 0$, solutions would converge a.e. to minima of W and that the interfaces between these sets (where the solution converges to minima) might develop a network structure that evolves under curve shortening flow. In [S1] was proved that this in fact holds for appropriate potentials and initial conditions if the considered networks are *triods* (networks that contain only three arcs). Moreover, it was proved that any smooth evolution of triods can be realized as a nodal set as described above. In order to prove Theorems 1.1 and 1.2 we show that such a representation can be extended to more general tree-like networks that have a regular evolution via network flow for $0 < t < T$.

We organize the paper as follows: In Section 2 we show that the topological class of network defines a unique coloring. In Section 3 we show that this coloring gives us an Allen-Cahn approximation. In Section 4 we use the previous approximation to conclude Theorems 1.1 and 1.2.

Remark 1.1. *We would like to remark that along the coming proofs all constants will be denoted by C, but they might vary from line to line.*

2. Topological class and coloring

Notice that any network $\mathcal{N} = \{\gamma_i\}$ contained in a set Ω defines a partition $\{\Omega_i\}$ of $\Omega \setminus \mathcal{N}$ as follows: $\partial\Omega_i \subset \mathcal{N}$, Ω_i is connected and $\bigcup_i \Omega_i = \Omega \setminus \mathcal{N}$ (see as an example Figure 2).

As stated in the introduction we are going to approximate solutions to the network flow via solutions to the Allen-Cahn equation with a three well potential. Let us assume that these three minima are given by c_1, c_2 and c_3. We will understand each of these minima as a color. It is expected that in each Ω_i the sequence of solutions u_ϵ to (1) converge to one of the c_i. Hence, a necessary requirement to approximate a network as an interface of the Allen-Cahn equation is that the partition defined by the network has a *three-coloring* as defined below:

Definition 2.1. *Suppose that we have a domain $\Omega \in \mathbb{R}^2$ and a partition $P = \{\Omega_i\}_{i=1}^n$ of Ω. Then we say that we can* three-color *P if for any region we can assign one of three fixed colors and the colors assigned in any adjacent regions are different.*

An example of a three-coloring would be in Figure 2 to associate c_1 to Ω_1 and Ω_4, c_2 to Ω_2 and c_3 to Ω_3 and Ω_5.

Suppose now that we have a tree-like graph that all its interior nodes are trivalent and the exterior ones are simple. Then the following proposition holds:

Proposition 2.1. *Suppose that we have a domain and a connected tree-like network with only trivalent nodes in the interior and simple nodes in the boundary. Let $P = \{\Omega_i\}$ be the partition associated to this network. There is a unique three-coloring of P (up to re-labeling*



FIGURE 2. Partition of the domain induced by a network

of the colors). Moreover, the same coloring can be associated to any other tree-like network that belongs to the same topological class.

Proof. For simplicity we may assume is included in a compact domain like in Figure 2 (otherwise the exterior vertices that we will refer to correspond to "vertices at infinity").

It is easy to see from the coming proof that the same coloring can be associated to any two networks that belong to the same topological class, hence we will not refer further to this assertion. For the existence and uniqueness we will use induction on the number of exterior vertices.

For $n = 3$ the result is trivial (each region has one of the color and this is unique, up to re-labeling of the colors). Suppose that the result holds for networks of trivalent interior nodes and $n-1$ simple exterior vertices. We prove that it also holds for networks of trivalent interior nodes and n simple exterior vertices. We will try to illustrate the induction procedure in Figures 3(a), 3(b) and 3(c) . Let us label the exterior vertices by $p_1, \dots, p_n$.

We first claim that there is an interior node v and two exterior nodes p_i and p_j such that there are edges $e_{p_i v}$ and $e_{p_j v}$ that connect p_i and v and p_j and v respectively. This can be easily proved by a combinatorial argument: Since the graph is a connected tree, we have that the number of interior nodes is equal to $n-2$ (this computation can be found for example in [MS]). Hence, it is not possible that each exterior node is connected to a separate interior node. Without loss of generality we can assume that the nodes adjacent to v are p_1 and p_2. And, since v is a trivalent node, there is only one remaining adjacent edge to v that we label by e.

Now remove p_1, v and the edge $e_{p_1 v}$ and we regard $e_{p_2 v} \bigcup e$ as one edge. Notice that by this procedure we eliminated only one region, which had $e_{p_1 v}$ and $e_{p_2 v}$ as boundaries. Moreover, we obtain a graph that topologically is equivalent to an tree-like graph with interior trivalent nodes and with $n-1$ single exterior nodes . By induction hypothesis we have that this graph can be uniquely colored with three colors (up to re-labeling of the colors). The edge that now is formed by $e_{p_2 v} \bigcup e$ has two adjacent regions that are colored by 2 different colors, let us say c_1 and c_2. Now we can add again p_1, v and the edge $e_{p_1 v}$. This procedure divides one of the regions above in two and adds one region, that can be colored by c_3. Since this region is only neighboring the regions that for the previous graph were adjacent to $e_{p_2 v} \bigcup e$, we have that this is a coloring of the graph with three colors.

To prove the uniqueness we just invert the procedure. Suppose that we have an arbitrary coloring of a network with n exterior nodes. We will show that this coloring agrees (up to



(a) The original tree-like network (b) Removing e_{p_1v} and using the induction hypothesis (c) adding e_{p_1v} and coloring R

FIGURE 3. Proof by induction of the existence of a coloring

re-labeling) with the one just constructed above. We can assume that p_1, p_2, v, e, e_{p_1v} and e_{p_2v} are defined as before. Via re-labeling we can color the region enclosed by e_{p_1v} and e_{p_2v} (that we label by R) with c_3. By removing e_{p_1v}, we have that this region becomes part of a region of either color c_1 or c_2. Let us assume it is c_1 and re-assign this color to region R (and the removed edge e_{p_1v}). Now, as before, we have a three-coloring of a tree-like graph with interior trivalent nodes and with $n-1$ single exterior nodes. By the uniqueness given by the induction hypothesis we conclude that (re-labeling, if necessary) the coloring for the network with the removed edge is the one used in the proof of existence above. Moreover, since the regions that were adjacent to R had color c_1 and c_2 the only possibly necessary rearrengement would be to interchange c_1 and c_2. Hence, following the existence proof, we conclude that the chosen arbitrary coloring (up to re-labeling) has to be the one constructed previously.

□

3. APPROXIMATION SKIM

In this section we prove that in fact a regular tree-like network can be understood as the nodal set of the limit of solutions to the Allen-Cahn. We assume that for positive times the nodes of the curves are trivalent and that the meeting angles of the curves at the nodes are all equal to $\frac{2\pi}{3}$. Although for certain networks it would be possible to extend the proof for other fixed meeting angle, in order to keep the presentation simpler, we will not discuss that situation. In our case it is enough to consider *symmetric* potentials W with three minima (where W attains the value 0). This corresponds to the following assumption $\Gamma(c_i, c_j) = \Gamma(c_j, c_k)$ for every $i \neq j$, $j \neq k$, where

$$\Gamma(\zeta_1,\zeta_2) = \inf\left\{ \int_0^1 W^{\frac{1}{2}}(\gamma(\lambda))|\gamma'(\lambda)|d\lambda : \gamma \in C^1([0,1],\mathbb{R}^2), \right. \\ \left. \gamma(0)=\zeta_1 \text{ and } \gamma(1)=\zeta_2 \right\}.$$

Moreover, we assume that there are three unique heteroclinics ζ_{ij} associated to this potential:*i.e* for each pair of minima c_i and c_j of W there is a unique curve ζ_{ij} that satisfies

$$\zeta_{ij}''(\lambda) + \frac{\nabla W(\zeta_{ij}(\lambda))}{2} = 0, \tag{4}$$

$$\lim_{\tau\to-\infty}\zeta_{ij}(\tau)=c_i,\quad \lim_{\tau\to\infty}\zeta_{ij}(\tau)=c_j, \tag{5}$$

In [S2] was proved that under this conditions, if W is regular enough, there is a solution $u_*(x):\mathbb{R}^2\to\mathbb{R}^2$ to

$$-\Delta u_*+\nabla W(u_*)=0,$$

which satisfies as $r\to\infty$

$$u_*(r\cos\theta,r\sin\theta)\to c_i \text{ for } \theta\in[\theta_{i-1},\theta_i], \tag{6}$$

$$u_*(r\cos\theta_i,r\sin\theta_i)\to\zeta_{ij}(0). \tag{7}$$

where the angles θ_i are given by the potential W in the following manner:

$$\frac{\sin\alpha_1}{\Gamma(c_2,c_3)}=\frac{\sin\alpha_2}{\Gamma(c_1,c_3)}=\frac{\sin\alpha_3}{\Gamma(c_1,c_2)},$$

with $\alpha_i=\theta_{i+1}-\theta_i$.

In [S1] it was proved:

Theorem 3.1. *Let $\mathcal{T}=\{\gamma^i\}$ be a triod evolving under curve shortening flow in a convex bounded domain Ω and let $O(t)$ be the point where the curves $\gamma_i(\cdot,t)$ meet. Assume that the angles at which the curves γ^i meet at the point $O(t)$ are fixed and that the lengths of the curves γ_i stay bounded away from 0. Consider a proper non-degenerate potential W with three global minima c_1,c_2 and c_3. Moreover, assume that this potential W is consistent with the fixed angles at $O(t)$. Then there are functions ϕ_ϵ, ψ_ϵ and v_ϵ such that there is a solution u_ϵ to (1)-(2)-(3) and*

$$\lim_{\epsilon\to0}v_\epsilon(x,t)\in\{c_1,c_2,c_3\}\ a.e., \tag{8}$$

$$\{(x,t):\lim_{\epsilon\to0}v_\epsilon(x,t)\notin\{c_1,c_2,c_3\}\}=\mathcal{T}\ \textit{and} \tag{9}$$

$$\lim_{\epsilon\to0}\sup_{\Omega\times[0,T]}|u_\epsilon-v_\epsilon|(x,t)=0. \tag{10}$$

There are two main ingredients on the proof of this theorem: the construction of the function v_ϵ and Lemma 3.1(that is stated below). In what follows we review these two elements of the proof.

We start by reviewing the construction of v_ϵ. Two regions can be distinguished: close to $O(t)$ and away from this point. In order to define these two regions, we consider $\tilde{\delta}$ (that needs to be chosen small enough) and we take a ball of radius $\tilde{\delta}$ around $O(t)$. Away from $O(t)$, namely on the complement of this ball we construct the function ϕ_ϵ as follows:

Let $d_i(x,t)=dist(x,\gamma^i(\cdot,t))$ the signed distance of a point $x\in\mathbb{R}^2$ to the curve $\gamma^i(\cdot,t)$ (where the signs of the distance functions need to be chosen appropriately) and consider $\tau^i(t)=(\cos\theta^i(t),\sin\theta^i(t))$ to be the unit tangents at $O(t)$. Define the sets:

$$D_{ii}(t)=\{x\in\Omega:d_i(x,t)\le\delta\}$$

$$D_{ii+1}(t)=\left\{x\in\Omega:d_i(x,t)\ge\frac{\delta}{2},\quad d_{i+1}(x,t)\le-\frac{\delta}{2}\text{ and }d_{ii+1}\ge\frac{\delta}{2}\right\}\subset S_{ii+1}.$$

Let $\xi_{ij}^{ext}(x,t)$ be a partition of unity associated to these sets, namely these functions satisfy $0 \le \xi_{ij}^{ext} \le 1$ (where $j \in \{i, i+1\}$), *supp* $\xi_{ij}^{ext} \subset D_{ij}$ (where *supp* denotes the support) and for every $x \in \bigcup_{i,j} D_{ij}$ holds $\sum_{i,j\in\{i,i+1\}} \xi_{ij}^{ext}(x,t) = 1$. Then for $x \in \bigcup_{i,j} D_{ij}$ we define

$$\phi_\epsilon(x,t) = \sum_{i=1}^{3} \left(\xi_{ii}^{ext}(x,t)\, \zeta_{ii+1}\left(\frac{d_i(x,t)}{\epsilon}\right) + \xi_{ii+1}^{ext}(x,t)\, c_i \right). \tag{11}$$

and $v_\epsilon(x,t) = \phi_\epsilon(x,t)$ for points x away from $O(t)$. Moreover, we use $\phi_\epsilon(x,t)$ as boundary condition in (3).

Furthermore, we extend the function ϕ_ϵ to the whole domain Ω as follows:

$$\phi_\epsilon^\eta(x,t) \equiv \left(1 - \eta_1\left(\frac{r(x,t)}{2\epsilon} + 1 - \frac{\tilde\delta}{2\epsilon}\right)\right) \phi_\epsilon(x,t),$$

where $r(x,t) = |x - O(t)|$ and $\eta_1 : \mathbb{R} \to \mathbb{R}$ is a function such that $\eta_1(x) \equiv 1$ when $|x| \le \frac{1}{2}$ and $\eta_1(x) \equiv 0$ for $|x| \ge 1$.

For points close to $O(t)$ we take $\frac{1}{2} < \rho < 1$ and define the regions $B_{\tilde\delta}(O(t)) \setminus B_{\epsilon^\rho}(O(t))$ and $B_{\frac{\epsilon^\rho}{2}}(O(t))$. Consider now $x \in B_{\tilde\delta}(O(t)) \setminus B_{\epsilon^\rho}(O(t))$ and take

$$\tilde\phi_\epsilon(x,t) = \sum_{i=1}^{3} \left(\xi_{2i}^{int}(\theta - \theta(t))\zeta_{ii+1}\left(\frac{d_i(x,t)}{\epsilon}\right) + \xi_{2i-1}^{int}(\theta - \theta(t)) c_i \right),$$

where $\theta(t)$ is the angle formed by the tangent $\tau^1(t)$ with the x-axis and $\{\xi_i^{int}\}_{i=1}^6$ is a partition of unity associated to the following family of intervals:

$$\mathcal{A}_{2i} = (\theta_i - \theta_{int}, \theta_i + \theta_{int})$$

$$\mathcal{A}_{2i+1} = \left(\theta_i + \frac{\theta_{int}}{2}, \theta_{i+1} - \frac{\theta_{int}}{2}\right),$$

with θ_{int} an angle chosen appropriately small and and θ_i the angles given by (6). Then define

$$\tilde\phi_\epsilon^\eta(x,t) \equiv \left(1 - \eta_2\left(\frac{x - O(t)}{\epsilon^\rho}\right)\right) \tilde\phi_\epsilon(x,t),$$

where $\eta_2 : \mathbb{R}^2 \to [0,1]$ is a function that satisfies $\eta_2(x) \equiv 1$ when $|x| \le \frac{1}{2}$.

On the other hand the function v_ϵ in the region that is ϵ^ρ close from the triple point is given by the stationary solution u_* to (1) that satisfies (6) and (7). More specifically, within the ball of radius $\tilde\delta$ we consider the function

$$\tilde v_\epsilon(x,t) = \tilde\phi_\epsilon^\eta(x,t) + \eta_2\left(\frac{x - O(t)}{\epsilon^\rho}\right) u_*\left(\frac{R_{\theta(t)}(x - O(t))}{\epsilon}\right), \tag{12}$$

where R_θ represents the rotation matrix by an angle θ.

Finally, we let

$$v_\epsilon(x,t) = \phi_\epsilon^\eta(x,t) + \eta_1\left(\frac{r(x,t)}{2\epsilon} + 1 - \frac{\tilde\delta}{2\epsilon}\right) \tilde v_\epsilon(x,t). \tag{13}$$

The initial condition to be chosen in (2) is

$$\psi_\epsilon(x) = v_\epsilon(x,0). \tag{14}$$

Before discussing the second main element of the proof, a few remarks are necessary. Let

$$F_\epsilon(h,\psi_\epsilon) = -\int_0^t \int_\Omega \mathcal{H}_\Omega(x,y,t-s)\left(\frac{\nabla_u W(h+\phi_\epsilon^\eta)}{\epsilon^2} + P\phi_\epsilon^\eta\right)(y,s)dyds$$

$$+\int_\Omega \mathcal{H}_\Omega(x,y,t)(\psi_\epsilon(y) - \phi_\epsilon^\eta(y,0))dy, \tag{15}$$

where $\mathcal{H}_\Omega$ denotes the heat kernel in Ω.

Notice that fixed points h_ϵ of this functional are solutions to the equation

$$\frac{\partial h_\epsilon}{\partial t} - \Delta h_\epsilon + \frac{\nabla_u W(h_\epsilon + \phi_\epsilon^\eta)}{2\epsilon^2} = -P\phi_\epsilon^\eta \quad \text{in } \Omega \tag{16}$$

$$h_\epsilon(x,t) = 0 \text{ on } \partial\Omega \tag{17}$$

$$h_\epsilon(x,0) = \psi(x) - \phi_\epsilon^\eta(x,0). \tag{18}$$

In particular, defining $u_\epsilon(x,t) = h_\epsilon(x,t) + \phi^\eta(x,t)$ we have $u_\epsilon(x,t)$ satisfies (1)-(2)-(3).

Now we can state the second main tool used to prove Theorem 3.1:

Lemma 3.1 (Lemma 4.1 in [S2]). *Fix $K > 0$. Consider the sequences of continuous functions ψ_n, w_n satisfying $\sup|\psi_n|$, $\sup|w_n| \leq K$. Let $\epsilon_n \to 0$ and $T_n > 0$. Assume in addition that for every $0 < \epsilon < 1$ holds $\sup_{x\in\Omega, t\in[0,T]} |h_\epsilon|(x,t) \leq K$. Then for each ψ_n, ϵ_n the functional F_{ϵ_n} has a unique fixed point h_{ϵ_n} and holds either*

1. $\lim_{n\to\infty} \sup_{\Omega\times[0,T_n]} |w_n - h_{\epsilon_n}| \to 0$, *or*
2. *there is a constant C, independent of ϵ_n and T_n such that*

$$\sup_{\Omega\times[0,T_n]} |w_n - h_{\epsilon_n}| \leq C \sup_{\Omega\times[0,T_n]} |F_{\epsilon_n}(w_n,\psi_n) - w_n|.$$

The proof of Theorem 3.1 follows from showing that $\sup_{\Omega\times[0,T_n]} |F_{\epsilon_n}(v_\epsilon,\psi_\epsilon) - v_\epsilon| \to 0$ as $\epsilon \to 0$ and the uniform bounds (independent of ϵ) of solutions to (1)-(2)-(3) proved in Lemma 2.2 of [S2].

Remark 3.1. *Suppose that there is an R such that $B_R \subset \Omega$. Then it is also possible in the definition of the functional F_ϵ (given by (15)) to consider instead of ϕ^η the function $\chi_R(x)\phi^\eta$ where $\chi_R : \bar{\Omega} \to [0,1]$ is a function that satisfies $\chi_R(x) = 0$ for $x \in B_{\frac{R}{2}}$ and $\chi_R(x) = 1$ for $x \in \partial\Omega$.*

In this case we would also need to substitute ϕ^η by $\chi_R(x)\phi^\eta$ in (16), (17) and (18). The proof of Lemma 3.1 nor the proof of Theorem 3.1 are altered by this modification.

Now we check that the proof of Theorem 3.1 extends for any tree-like network. More specifically we show that

Theorem 3.2. *Let $\mathcal{N} = \{\gamma^i\}_{i=1}^{2n-3}$ be a network evolving under curve shortening flow in a convex bounded domain Ω at times $t \in [0,T)$ and let $\{O_i(t)\}_{i=1}^{n-2}$ be its interior nodes. Assume that the evolution is regular for all $0 < t \leq T$ and that the lengths of the curves γ_i stay bounded away from 0. Moreover, we assume that the nodes stay a fixed distance away from the boundary and from each other and that their linear and angular speeds remain bounded. The curvature of the arcs is assumed to be bounded by $\frac{C}{\sqrt{t}}$ where C is a fixed constant.*

Consider a proper non-degenerate symmetric potential W with three global minima c_1, c_2 and c_3. Then there are functions ϕ_ϵ, ψ_ϵ and n_ϵ such that there is a solution u_ϵ to (1)-(2)-(3) that satisfies

$$\lim_{\epsilon\to 0} n_\epsilon(x,t) \in \{c_1, c_2, c_3\} \text{ a.e.,} \tag{19}$$

$$\{(x,t) : \lim_{\epsilon\to 0} n_\epsilon(x,t) \notin \{c_1, c_2, c_3\}\} = \mathcal{N} \text{ and} \tag{20}$$

$$\lim_{\epsilon\to 0} \sup_{\Omega\times[0,T]} |u_\epsilon - n_\epsilon|(x,t) = 0. \tag{21}$$

Furhermore, $\lim_{\epsilon\to 0} n_\epsilon(x,t)$ defines a coloring of the network $\mathcal{N}$.

We will only briefly sketch the proof of this theorem. For further details on the missing computations we refer the reader to [S1].

Proof. The proof of this theorem is completely analogous to the proof of Theorem 3.1.

We start by constructing the function n_ϵ: We consider a coloring of the partition associated to the network and define the function $col : \Omega \setminus \mathcal{N}(\cdot,t) \times [0,T) \to \{c_i\}_{i=1}^3$ as $col(x,t) = c_i$ if the color associated to x at time t is c_i. Notice that as long as the curves do no not disappear during the evolution, the topological class will not change, hence, the coloring is in fact preserved in time.

As before, we denote $d_i(x,t) = dist(x, \gamma^i(\cdot,t))$ the signed distance of a point $x \in \mathbb{R}^2$ to the curve $\gamma^i(\cdot,t)$, where the signs of the distance functions need to be chosen appropriately.

Analogous to the sets D_{ij} defined in the proof of Theorem 3.1, we consider:

$$\mathcal{D}_i(t) = \{x \in \Omega : d_i(x,t) \le \delta\}$$

$$\mathcal{C}_i(t) = \left\{ x \in \Omega : col(x,t) = c_i \text{ and } |d_j(x,t)| \ge \frac{\delta}{2} \text{ for every } j \right\}.$$

Define a partition of unity $\{\xi^{ext}_{\mathcal{D}_j}, \quad \xi^{ext}_{\mathcal{C}_j}\}$, associated to these sets. We denote $Z_{\mathcal{D}_i}(\cdot) = \zeta_{jk}(\cdot)$ if $\gamma_i \in \partial\mathcal{C}_j \bigcap \partial\mathcal{C}_k$.

Then we can define for $x \in \bigcup_j \mathcal{D}_j \cup \bigcup_j \mathcal{C}_j$ the function

$$\phi_\epsilon(x,t) = \sum_i \xi^{ext}_{\mathcal{D}_i}(x,t)\, Z_{\mathcal{D}_i}\left(\frac{d_i(x,t)}{\epsilon}\right) + \sum_i \xi^{ext}_{\mathcal{C}_i}(x,t)\, col(x,t). \tag{22}$$

Here we would like to remark that the appropriate choice of sign of d_i corresponds to have for $x \in \mathcal{C}_j(t)$ that $Z_{\mathcal{D}_i}\left(\frac{d_i(x,t)}{\epsilon}\right) \to c_j$ as $\epsilon \to 0$.

This function will correspond to the function n_ϵ away from the nodes. We also use this function as boundary condition in (3).

Now we extend the function ϕ_ϵ to the whole domain Ω using appropriate cut-off functions. Namely we take

$$\phi^\eta_\epsilon(x,t) \equiv \left(\sum_{i=1}^{n-2} \left(1 - \eta_1 \left(\frac{r_i(x,t)}{2\epsilon} + 1 - \frac{\tilde{\delta}}{2\epsilon} \right) \right) \right) \phi_\epsilon(x,t),$$

where $r_i(x,t) = |x - O_i(t)|$ and $\eta_1 : \mathbb{R} \to \mathbb{R}$ is a function that satisfies $\eta_1(x) \equiv 1$ when $|x| \le \frac{1}{2}$ and $\eta_1(x) \equiv 0$ for $|x| \ge 1$.

Now we extend the construction close to the nodes $O_i(t)$. Since we have a coloring, around each node $O_i(t)$ we can do a construction analogous to (12) and consistent with the coloring. Let $\tau_i^j(t) = (\cos\theta_i^j(t), \sin\theta_i^j(t))$ to be the unit tangents at $O_i(t)$. In the region ϵ^ρ far from the triple point we consider a partition of unity $\{\xi_i^{int}\}_{i=1}^3$ associated to the family of intervals $\{\mathcal{A}_j\}_{j=1}^6$, where

$$\mathcal{A}_{2j} = \left(\frac{2j\pi}{3} - \theta_{int}, \frac{2j\pi}{3} + \theta_{int}\right)$$

$$\mathcal{A}_{2j+1} = \left(\frac{2j\pi}{3} + \frac{\theta_{int}}{2}, \frac{2(j+1)\pi}{3} - \frac{\theta_{int}}{2}\right),$$

and θ_{int} is an angle chosen appropriately small. We define $Z_{\mathcal{A}_{2i}}(\cdot) = \zeta_{jk}(\cdot)$ if $col(x,t) = c_j$ for $x \in \mathcal{A}_{2i+1}$ and $col(x,t) = c_k$ for $x \in \mathcal{A}_{2i-1}$. We label the adjacent arcs to $O_i(t)$ by $\{\gamma_i^j\}_{j=1}^3$ and the corresponding distance $d_i^j(x,t) = dist(x, \gamma_i^j(\cdot,t))$. Moreover, we assume that $\gamma_i^j \bigcap \mathcal{A}_{2j} \neq \emptyset$ (while $\gamma_i^j \bigcap \mathcal{A}_{2k} = \emptyset$ for every $k \neq 2j$). Then we take the function

$$\tilde{\phi}_\epsilon^i(x,t) = \sum_{j=1}^3 \left(\xi_{2j}^{int}(\theta - \theta^i(t)) Z_{\mathcal{A}_{2i}}\left(\frac{d_i^j(x,t)}{\epsilon}\right) + \xi_{2i-1}^{int}(\theta - \theta(t)) col(x,t)\right),$$

where $\theta^i(t)$ is the angle formed by the tangent $\tau_1^i(t)$ with the x-axis. As before we extend the function $\tilde{\phi}_\epsilon^i$ to the whole domain by multiplying by an appropriate cut-off function:

$$\tilde{\tilde{\phi}}_\epsilon^i(x,t) \equiv \left(1 - \eta_2\left(\frac{x - O_i(t)}{\epsilon^\rho}\right)\right) \tilde{\phi}_\epsilon^i(x,t),$$

with $\frac{1}{2} < \rho < 1$ as before and $\eta_2 : \mathbb{R}^2 \to \mathbb{R}$ is a smooth function such that $\eta_2(x) \equiv 1$ when $|x| \leq \frac{1}{2}$.

The function n_ϵ in the region that is ϵ^ρ close from the triple point is given by the stationary solution u_* to (1) that satisfies (6) and (7). More precisely we take

$$\tilde{n}_\epsilon^i(x,t) = \tilde{\tilde{\phi}}_\epsilon^i(x,t) + \eta_2\left(\frac{x - O_i(t)}{\epsilon^\rho}\right) u_*\left(\frac{R_{\theta^i(t)}(x - O_i(t))}{\epsilon}\right), \tag{23}$$

where R_θ represents the rotation matrix by an angle θ.

Finally we define

$$n_\epsilon(x,t) = \phi_\epsilon^\eta(x,t) + \sum_i \eta_1\left(\frac{r_i(x,t)}{2\epsilon} + 1 - \frac{\tilde{\delta}}{2\epsilon}\right) \tilde{n}_\epsilon^i(x,t). \tag{24}$$

The initial condition (2) is given by:

$$\psi_\epsilon(x) = n_\epsilon(x,0). \tag{25}$$

It is easy to see that the computations in [S1] (used to show that $|F_{\epsilon_n}(v_\epsilon, \psi_\epsilon) - v_\epsilon| \to 0$ uniformly as $\epsilon \to 0$) can be extended to the function n_ϵ constructed above. We will not review this computation here. However an analogous computation will be presented in the coming proof .

Now, as in [S1], we can use Lemma 3.1 and the uniform bounds provided by Lemma 2.2 in [S2] to conclude the result. □

Moreover, we extend this result to the following situation

Theorem 3.3. *Let* $\mathcal{N} = \{\gamma^i\}_{i=1}^{2n-3}$ *be a network evolving under curve shortening flow in a convex bounded domain* Ω *and let* $\{O_i(t)\}_{i=1}^{n-2}$ *be its interior nodes. Assume that the angles at which the curves* γ^i *meet at the points* $O_i(t)$ *are of* $\frac{2\pi}{3}$ *and that there is a constant* C *such that the lengths of the curves* γ_i *and the distances between the interior nodes are bounded below by* $C\sqrt{t}$*. Moreover, we assume that the nodes stay a fixed distance away form the boundary and that their linear and angular speeds are uniformly bounded. The curvature of the arcs is assumed to be bounded by* $\frac{C}{\sqrt{t}}$ *where* C *is a fixed constant. Consider a proper non-degenerate symmetric potential* W *with three global minima* c_1, c_2 *and* c_3*.Then there are functions* ϕ_ϵ*,* ψ_ϵ *and* n_ϵ *such that there is a solution* u_ϵ *to (1)-(2)-(3) and*

$$\lim_{\epsilon\to 0} n_\epsilon(x,t) \in \{c_1, c_2, c_3\} \text{ a.e.,} \tag{26}$$

$$\{(x,t) : \lim_{\epsilon\to 0} n_\epsilon(x,t) \notin \{c_1, c_2, c_3\}\} = \mathcal{N} \text{ and} \tag{27}$$

$$\lim_{\epsilon\to 0} \sup_{\Omega} |u_\epsilon - n_\epsilon|(\cdot, t) = 0 \text{ for every } t \in [0,T]. \tag{28}$$

Furthermore, $\lim_{\epsilon\to 0} n_\epsilon(x,t)$ *defines a coloring of the network* $\mathcal{N}$*.*

Before proving this Theorem we will need a weaker version of Lemma 3.1:

Lemma 3.2. *Consider the functional* F_ϵ *in Lemma 3.1. Fix* $K > 0$*. Consider the sequences of continuous functions* ψ_n, w_n *satisfying* $\sup|\psi_n|,\ \sup|w_n| \le K$*. Let* $\epsilon_n \to 0$ *and* $0 < T_n \le T$*. Assume in addition that for every* $0 < \epsilon < 1$ *holds* $\sup_{x\in\Omega, t\in[0,T]} |h_\epsilon|(x,t) \le K$*. Then for each* ψ_n, ϵ_n *the functional* F_{ϵ_n} *has a unique fixed point* h_{ϵ_n} *and holds either*

1. $\lim_{n\to\infty} \sup_{\Omega\times[0,T_n]} t|w_n - h_{\epsilon_n}| \to 0$*, or*
2. *there is a constant* C*, independent of* ϵ_n *and* T_n *such that*

$$\sup_{\Omega\times[0,T_n]} t|w_n - h_{\epsilon_n}| \le C \sup_{\Omega\times[0,T_n]} t|F_{\epsilon_n}(w_n,\psi_n) - w_n|.$$

Proof. Let us assume that neither (1) nor (2) hold. Then for every n there are T_n, ϵ_n and w_n such that

$$\sup_{\Omega\times[0,T_n]} t|w_n - h_{\epsilon_n}| \ge n \sup_{\Omega\times[0,T_n]} t|F_{\epsilon_n}(w_n,\psi_n) - w_n|.$$

Fix $\delta > 0$. Since w_n and h_{ϵ_n} are uniformly bounded, we have $\sup_{\Omega\times[\delta,T_n]} |F_{\epsilon_n}(w_n,\psi_n) - w_n| \frac{1}{\delta} \sup_{\Omega\times[\delta,T_n]} t|F_{\epsilon_n}(w_n,\psi_n) - w_n| \to 0$. From Lemma 3.1 we have that

$$\lim_{n\to\infty} \sup_{\Omega\times[\delta,T_n]} |w_n - h_{\epsilon_n}| \to 0.$$

Hence

$$\begin{aligned}\lim_{n\to\infty} \sup_{\Omega\times[0,T_n]} t|w_n - h_{\epsilon_n}| &\le \sup\{T \sup_{\Omega\times[\delta,T_n]} |w_n - h_{\epsilon_n}|,\ \sup_{\Omega\times[0,\delta]} t|w_n - h_{\epsilon_n}|\} \\ &\le \sup_{\Omega\times[0,\delta]} t|w_n - h_{\epsilon_n}| \\ &\le C\delta.\end{aligned}$$

Since δ is arbitrary we conclude that $\lim_{n\to\infty} \sup_{\Omega\times[0,T_n]} t|w_n - h_{\epsilon_n}| = 0$, which contradicts that (1) does not hold.

□

Now we can prove Theorem 3.3:

Proof of Theorem 3.3. The proof of this Theorem is analogous to the one of the previous one. However, some modifications both in the construction of the function n_ϵ and in the computation of the bounds that we use to conclude are necessary.

We would first like to remark that the hypotheses of the theorem allow that some of the vertices at time $t = 0$ split into several for positive times. However, in the construction below we would like to have the same number of nodes for all times. Hence, the considered nodes $\{O_i(t)\}_{i=1}^{n-2}$ will be the ones that are different for positive times and some of them might agree for t=0. For example, for the self-similar solutions coming out of n half-lines that start at the origin, we have $n-2$ nodes $\{O_i(t)\}_{i=1}^{n-2}$ that agree at $t = 0$.

We start by discussing the function n_ϵ that will be used. The notation will be as in the previous proof. We first modify the sets $\mathcal{D}_i$ and $\mathcal{C}_i$ as follows:

$$\mathcal{D}_i^\epsilon(t) = \{x \in \Omega : d_i(x,t) \le \frac{\epsilon^\rho}{2}\}$$

$$\mathcal{C}_i^\epsilon(t) = \left\{ x \in \Omega : col(x,t) = c_i \text{ and } |d_j(x,t)| \ge \frac{\epsilon^\rho}{4} \text{ for every } j \right\}.$$

Now $\{\xi^{ext}_{\mathcal{D}_j^\epsilon},\ \ \xi^{ext}_{\mathcal{C}_j^\epsilon}\}$ denotes a partition of unity associated to these sets. As before, we define $Z_{\mathcal{D}_i^\epsilon}(\cdot) = \zeta_{jk}(\cdot)$ if $\gamma_i \in \partial\mathcal{C}_j^\epsilon \bigcap \partial\mathcal{C}_k^\epsilon$.

Then we define the function ϕ_ϵ for $x \in \bigcup_j \mathcal{D}_j^\epsilon \cup \bigcup_j \mathcal{C}_j^\epsilon$ (that will be used as the new boundary condition in (3)) by

$$\phi_\epsilon(x,t) = \sum_i \xi^{ext}_{\mathcal{D}_i^\epsilon}(x,t)\, Z^\epsilon_{\mathcal{D}_i}\left(\frac{d_i(x,t)}{\epsilon}\right) + \sum_i \xi^{ext}_{\mathcal{C}_i^\epsilon}(x,t)\, col(x,t). \tag{29}$$

The signs for d_i are chosen appropriately as before and the following extension for ϕ_ϵ are considered:

$$\phi_\epsilon^\eta(x,t) \equiv \left(\sum_{i=1}^{n-2}\left(1 - \eta_2\left(\frac{x - O_i(t)}{\epsilon^\rho}\right)\right)\right)\phi_\epsilon(x,t),$$

where $\eta_2 : \mathbb{R} \to \mathbb{R}$ is the cut-off function described in previous proof. Now we define

$$n_\epsilon(x,t) = \phi_\epsilon^\eta(x,t) + \sum_{i=1}^{n-2} \eta_2\left(\frac{x - O_i(t)}{\epsilon^\rho}\right) u_*\left(\frac{R_{\theta^i(t)}(x - O_i(t))}{\epsilon}\right) \tag{30}$$

and the initial condition (2) is given by:

$$\psi_\epsilon(x) = n_\epsilon(x,0). \tag{31}$$

Let $w_\epsilon(x,t) = n_\epsilon(x,t) - \phi_\epsilon^\eta(x,t)$. Correspondingly, if we define the functional as in Remark 3.1, we would take $w_\epsilon(x,t) = n_\epsilon(x,t) - \xi_R(x)\phi_\epsilon^\eta(x,t)$. If the latter choice were made, in all the computations below the function ϕ_ϵ^η would need to be replaced by $\xi_R(x)\phi_\epsilon^\eta(x,t)$ and the same conclusions would hold. We leave this case to be checked by the reader.

Since $w_\epsilon(x,t) = 0$ for $x \in \partial\Omega$, we can write

$$w_\epsilon(x,t) = \int_0^t \int_\Omega \mathcal{H}_\Omega(x,y,t-s) P w_\epsilon(y,s)\,dyds + \int_\Omega \mathcal{H}_\Omega(x,y,t) w_\epsilon(y,0)\,dy.$$

Which implies

$$
\begin{aligned}
(F_\epsilon(w_\epsilon, \psi_\epsilon) - w_\epsilon)(x,t) &= \int_0^t \int_\Omega \mathcal{H}_\Omega(x,y,t-s) \left(-\frac{\nabla_u W(w_\epsilon + \phi_\epsilon^\eta)}{\epsilon^2}(y,s) - P(\phi_\epsilon^\eta + w_\epsilon)(y,s) \right) dyds \\
&\quad + \int_\Omega \mathcal{H}_\Omega(x,y,t)(\psi_\epsilon(y) - \phi_\epsilon^\eta(y) - w_\epsilon(y,0))dy \\
&= \int_0^t \int_\Omega \mathcal{H}_\Omega(x,y,t-s) \left(-\frac{\nabla_u W(n_\epsilon)}{\epsilon^2} - Pn_\epsilon \right)(y,s)dyds. \qquad (32)
\end{aligned}
$$

Hence, it is enough to prove that

$$
\sup_{\Omega\times[0,T)} t \left| \int_0^t \int_\Omega \mathcal{H}_\Omega(x,y,t-s) \left(\frac{\nabla_u W(n_\epsilon)}{\epsilon^2} + Pn_\epsilon \right)(y,s)dyds \right| \to 0 \text{ as } \epsilon \to 0.
$$

We start by briefly reviewing the computation away from the nodes. This computation, for all times, is analogous to the ones used to prove Theorem 3.1 in [S1]. We bound:

$$
I_\epsilon^{ext}(x,t) = t \left| \int_0^t \int_{\Omega\setminus\bigcup_i B_{\epsilon^\rho}(O_i(s))} \mathcal{H}_\Omega(x,y,t-s) \left(\frac{\nabla_u W(n_\epsilon)}{\epsilon^2} + Pn_\epsilon \right)(y,s)dyds \right|
$$

for any $x \in \Omega$ and $0 \le t \le T$. This integral can be subdivided as $I_\epsilon^{ext} \le I^{ext}_{\mathcal{D}_i^\epsilon \setminus (\mathcal{D}_i^\epsilon \bigcap \mathcal{C}_i^\epsilon)} + I^{ext}_{\mathcal{C}_i^\epsilon \setminus (\mathcal{D}_i^\epsilon \bigcap \mathcal{C}_i^\epsilon)} + I^{ext}_{\mathcal{D}_i^\epsilon \bigcap \mathcal{C}_i^\epsilon}$, where

$$
\begin{aligned}
A(x,y,t,s) =& \mathcal{H}_\Omega(x,y,t-s) \left(\frac{\nabla_u W(n_\epsilon)}{\epsilon^2} + Pn_\epsilon \right)(y,s), \\
I^{ext}_{\mathcal{D}_i^\epsilon \setminus (\mathcal{D}_i^\epsilon \bigcap \mathcal{C}_i^\epsilon)} =& t \left| \int_0^t \int_{\mathcal{D}_i^\epsilon \setminus (\mathcal{D}_i^\epsilon \bigcap \mathcal{C}_i^\epsilon)} A(x,y,t,s)dyds \right| \\
I^{ext}_{\mathcal{C}_i^\epsilon \setminus (\mathcal{D}_i^\epsilon \bigcap \mathcal{C}_i^\epsilon)} =& y \left| \int_0^t \int_{\mathcal{C}_i^\epsilon \setminus (\mathcal{D}_i^\epsilon \bigcap \mathcal{C}_i^\epsilon)} A(x,y,t,s)dyds \right| \\
I^{ext}_{\mathcal{D}_i^\epsilon \bigcap \mathcal{C}_i^\epsilon} =& t \left| \int_0^t \int_{\mathcal{D}_i^\epsilon \bigcap \mathcal{C}_i^\epsilon} A(x,y,t,s)dyds \right|
\end{aligned}
$$

Using the definition of n_ϵ it is not hard to compute that

$$
\begin{aligned}
|A(x,y,t,s)| =& \quad \left| \mathcal{H}_\Omega(x,y,t-s) \tfrac{k_i^2(\lambda,t)d_i}{\epsilon(1+k_i(\lambda,t)d_i)} \zeta'_{ii+1} \right| \\
\le& \quad \mathcal{H}_\Omega(x,y,t-s) \left| \tfrac{k_i^2(\lambda,t)d_i}{\epsilon(1+k_i(\lambda,t)d_i)} \right| e^{-c\frac{d_i(x,t)}{\epsilon}} && \text{for } x \in \mathcal{D}_i^\epsilon \setminus (\mathcal{D}_i^\epsilon \bigcap \mathcal{C}_i^\epsilon). \\
|A(x,y,t,s)| =& \quad 0 && \text{for } x \in \mathcal{C}_i^\epsilon \setminus (\mathcal{D}_i^\epsilon \bigcap \mathcal{C}_i^\epsilon). \\
|A(x,y,t,s)| =& \quad |\mathcal{H}_\Omega(x,y,t-s)| \left| Pn_\epsilon + \tfrac{\nabla_u W(n_\epsilon)}{\epsilon^2} \right| \\
& \quad \le C \max\left\{ \sup_i |k_i|, \tfrac{e^{-c\epsilon^{\rho-1}}}{\epsilon^2} \right\} && \text{for } x \in \mathcal{D}_i^\epsilon \bigcap \mathcal{C}_i^\epsilon. \\
\left| Pn_\epsilon + \tfrac{\nabla_u W(n_\epsilon)}{\epsilon^2} \right| & \quad \to 0 \text{ as } \epsilon \to 0 && \text{for } x \in \mathcal{D}_i^\epsilon \bigcup \mathcal{C}_i^\epsilon.
\end{aligned}
$$

Since k^i is an integrable function, the Dominated Convergence Theorem implies that

$$
\lim_{\epsilon\to 0} I_\epsilon^{ext} = 0. \qquad (33)
$$

For further details in this computation we refer the reader to [S1].

Now we need to bound

$$I_\epsilon^{int}(x,t) = t\left|\int_0^t\int_{\bigcup_i B_{\epsilon^\rho}(O_i(s))}\mathcal{H}_\Omega(x,y,t-s)\left(\frac{\nabla_u W(n_\epsilon)}{\epsilon^2}+Pn_\epsilon\right)(y,s)dyds\right|.$$

Notice that there is a K that depends only on the network such that for $t \geq K\epsilon^{2\rho}$ all the nodes are already separated enough to make the definition of n_ϵ appropriate to distinguish each region. It is not hard to check, that the computations in [S1] can be easily extended for these large times. However, since at $t=0$ there might be nodes that coincide (such as in the case of the self-similar solutions described in the introduction), for $t \leq K\epsilon^{2\rho}$ the computations need to be handled more carefully near this multiple nodes. In order to have some room in our calculations, we will divide the times into slightly larger regions, namely $t \leq \epsilon^\rho$ and $t \geq \epsilon^\rho$. We will first focus on bounding $I_\epsilon^{int}(x,t)$ for small times.

Since the nodes $O_i(t)$ are away form the boundary, we have

$$\mathcal{H}_\Omega(x,y,t-s)\left|\frac{\nabla_u W(n_\epsilon)}{\epsilon^2}+Pn_\epsilon\right| \leq \frac{C}{\epsilon^2}\frac{e^{-\frac{|x-y|^2}{t-s}}}{t-s}.$$

Then it follows for $t \leq \epsilon^\rho$:

$$\begin{aligned}
I_\epsilon^{int}(x,t) \leq & \frac{C}{\epsilon^2}t\int_0^t\int_{\bigcup_i B_{\epsilon^\rho}(O_i(s))}\frac{e^{-\frac{|x-y|^2}{t-s}}}{t-s}dyds\\
\leq & \frac{C\epsilon^\rho}{\epsilon^2}\int_0^{t-\epsilon^3}\int_{B_{\epsilon^\rho}}\frac{1}{t-s}dyds, + \int_{t-\epsilon^3}^t\int_{\mathbb{R}^2}\frac{e^{-\frac{|x-y|^2}{t-s}}}{t-s}dyds\\
\leq & \frac{C\epsilon^\rho}{\epsilon^2}\left(\pi\epsilon^{2\rho}(\ln t - 3\ln\epsilon)+\epsilon^3\right)\\
\leq & -C\epsilon^{3\rho-2}\ln\epsilon
\end{aligned}$$

Hence we have,

$$I_\epsilon^{int}(x,t) \leq -C\epsilon^{3\rho-2}\ln\epsilon, \text{ for } t\in[0,\epsilon^\rho]. \tag{34}$$

Consequently, we choose $\frac{1}{2} < \frac{2}{3} < \rho < 1$.

For $t \geq \epsilon^\rho$ we bound the integral as follows:

$$I_\epsilon^{int} \leq T(I_{s\leq K\epsilon^{2\rho}}^{int} + I_{s\geq K\epsilon^{2\rho}}^{int} + I_{s\geq K\epsilon^{2\rho}}^{trans}),$$

where

$$\begin{aligned}
I_{s\leq K\epsilon^{2\rho}}^{int} &= \left|\int_0^{K\epsilon^{2\rho}}\int_{\bigcup_i B_{\epsilon^\rho}(O_i(s))}\mathcal{H}_\Omega(x,y,t-s)\left(\frac{\nabla_u W(n_\epsilon)}{\epsilon^2}+Pn_\epsilon\right)(y,s)dyds\right|\\
I_{s\geq K\epsilon^{2\rho}}^{int} &= \left|\int_{K\epsilon^{2\rho}}^t\int_{\bigcup_i B_{\frac{\epsilon^\rho}{2}}(O_i(s))}\mathcal{H}_\Omega(x,y,t-s)\left(\frac{\nabla_u W(n_\epsilon)}{\epsilon^2}+Pn_\epsilon\right)(y,s)dyds\right|\\
I_{s\geq K\epsilon^{2\rho}}^{trans} &= \left|\int_{K\epsilon^{2\rho}}^t\int_{\bigcup_i B_{\epsilon^\rho}(O_i(s))\setminus B_{\frac{\epsilon^\rho}{2}}(O_i(s))}\mathcal{H}_\Omega(x,y,t-s)\left(\frac{\nabla_u W(n_\epsilon)}{\epsilon^2}+Pn_\epsilon\right)(y,s)dyds\right|.
\end{aligned}$$

For $s \leq K\epsilon^{2\rho}$, as in the computation for (34), we have

$$\mathcal{H}_\Omega(x,y,t-s)\left|\frac{\nabla_u W(n_\epsilon)}{\epsilon^2} + Pn_\epsilon\right|(y,s) \leq \frac{C}{\epsilon^2}\frac{e^{-\frac{|x-y|^2}{t-s}}}{t-s}.$$

Then

$$\begin{aligned}
I^{int}_{s\leq K\epsilon^{2\rho}}(x,t) \leq & \frac{C}{\epsilon^2}\int_0^{K\epsilon^{2\rho}}\int_{\bigcup_i B_{\epsilon^\rho}(O_i(s))}\frac{e^{-\frac{|x-y|^2}{t-s}}}{t-s}dyds\\
\leq & \frac{C}{\epsilon^2}\left(\int_0^{K\epsilon^{2\rho}}\int_{\bigcup_i B_{\epsilon^\rho}(O_i(s))}\frac{1}{t-s}dyds\right)\\
\leq & -\frac{C}{\epsilon^2}\pi\epsilon^{2\rho}\sum_i \ln(1-\frac{K\epsilon^{2\rho}}{t})\\
\leq & -C\frac{\sum_i \ln(1-K\epsilon^\rho)}{\epsilon^{2(1-\rho)}}\\
\leq & C(n-2)\epsilon^{3\rho-2}.
\end{aligned}$$

Since we took $\frac{1}{2} < \frac{2}{3} < \rho < 1$, we have

$$I^{int}_{s\leq K\epsilon^{2\rho}}(x,t) \leq C\epsilon^{\tilde{\rho}} \text{ for some } \tilde{\rho} > 0 \tag{35}$$

For $I^{int}_{s\geq K\epsilon^{2\rho}}$ we proceed as in [S1]. It is easy to compute that

$$\begin{aligned}
|A(x,y,t,s)| = & \left|\mathcal{H}_\Omega(x,y,t-s)Ju_*\frac{\theta' R_{\theta(t)+\frac{\pi}{2}}(x-O(t)) - R_{\theta(t)}O'(t)}{\epsilon}\right|\\
\leq & \frac{C}{\epsilon}\mathcal{H}_\Omega(x,y,t-s) \text{ for } y \in B_{\frac{\epsilon^\rho}{2}}(O_i(t))
\end{aligned}$$

where, as before, $A(x,y,t,s) = \mathcal{H}_\Omega(x,y,t-s)\left(\frac{\nabla_u W(n_\epsilon)}{\epsilon^2} + Pn_\epsilon\right)(y,s)$.

Since the nodes are away from the boundary, we have that $|\mathcal{H}_\Omega(x,y,t-s)| \leq \frac{Ce^{-\frac{|x-y|^2}{t-s}}}{t-s}$ and

$$\begin{aligned}
I^{int}_{s\geq K\epsilon^{2\rho}} \leq & \frac{C}{\epsilon}\int_{K\epsilon^{2\rho}}^{t}\int_{B_{\frac{\epsilon^\rho}{2}}(O(s))}\frac{e^{-\frac{|x-y|^2}{t-s}}}{t-s}dyds\\
\leq & \frac{C}{\epsilon}\left(\int_{K\epsilon^{2\rho}}^{t-\epsilon^m}\int_{B_{\frac{\epsilon^\rho}{2}}(O(s))}\frac{1}{t-s}dyds + \int_{t-\epsilon^m}^{t}\int_{\mathbb{R}^2}\frac{e^{-\frac{|x-y|^2}{t-s}}}{t-s}dyds\right)\\
\leq & \frac{C}{\epsilon}\left(\int_{K\epsilon^{2\rho}}^{t-\epsilon^m}\frac{\pi\epsilon^{2\rho}}{4(t-s)}ds + \int_{t-\epsilon^m}^{t}ds\right)\\
\leq & \frac{C}{\epsilon}\left((\ln(t-K\epsilon^{2\rho}) - m\ln\epsilon)\frac{\pi\epsilon^{2\rho}}{4} + \epsilon^m\right)\\
\leq & C\left((\ln T - m\ln\epsilon)\frac{\pi\epsilon^{2\rho-1}}{4} + \epsilon^{m-1}\right) \to 0 \text{ as } \epsilon \to 0.
\end{aligned} \tag{36}$$

For $t \geq \epsilon^\rho$ in $B_{\epsilon^\rho}(O_i(t)) \setminus B_{\frac{\epsilon^\rho}{2}}(O_i(t))$ we have that $\eta_2\left(\frac{x-O_j(t)}{\epsilon^\rho}\right) = 0$ and $n_\epsilon = \phi_\epsilon(x,t) + \eta_2\left(\frac{x-O_i(t)}{\epsilon^\rho}\right)\left(u_*\left(\frac{R_{\theta^i(t)}(x-O_i(t))}{\epsilon}\right) - \phi_\epsilon(x,t)\right)$. Hence

$$\begin{aligned} Pv_\epsilon + \frac{\nabla_u W(v_\epsilon)}{\epsilon^2} &= \left(P\phi_\epsilon(x,t) + \frac{\nabla_u W(\phi_\epsilon)}{\epsilon^2}\right) \\ &\quad + P\left[\eta_2\left(\frac{x-O_i(t)}{\epsilon^\rho}\right)\left(u_*\left(\frac{R_{\theta^i(t)}(x-O_i(t))}{\epsilon}\right) - \phi_\epsilon(x,t)\right)\right] \\ &\quad + \frac{\nabla_u W(v_\epsilon)}{\epsilon^2} - \frac{\nabla_u W(\phi_\epsilon)}{\epsilon^2}. \end{aligned} \tag{37}$$

The first parenthesis can be bounded as (33), while the bounds for the other quantities follow from the fast convergence of u_* to ϕ_ϵ (see [S1] and [S2] for details on this). Hence we have for any $m > 0$

$$I^{trans}_{s \geq K\epsilon^{2\rho}} \leq C\left(\int_0^t \int_{B_{\epsilon^\rho}(O(t)) \setminus B_{\frac{\epsilon^\rho}{2}}(O(t))} \mathcal{H}_\Omega(x,y,t-s)\left|P\phi_\epsilon + \frac{\nabla_u W(\phi_\epsilon)}{\epsilon^2}\right|(y,s) + \epsilon^m\right). \tag{38}$$

We conclude from Lemma 3.2 and (33), (34), (35), (36) and (38) that for every $t > 0$ the result holds. However, since the initial condition was appropriately chosen, we in fact have that the result holds for every $t \geq 0$.

□

Remark 3.2. *Notice that the self-similar solutions to the network flow studied in* [MS] *satisfy the conditions of Theorem 3.3 in a large ball for a fixed time (that depends on the size of the domain). The boundary condition needs to be chosen as the motion of the intersection point of the solution and the boundary of the ball.*

4. Proof of Theorems 1.1 and 1.2

Theorems 1.1 is a particular case of Theorem 1.2 . Hence, we only treat the more general case. Suppose that we have two networks $\mathcal{N}_1$ and $\mathcal{N}_2$ in a domain Ω. Furthermore assume that they have the same topological type and their initial conditions coincide.

Since Theorem 3.3 was proved in a bounded domain, if Ω is unbounded we define the bounded domain Ω_R as the intersection of a large ball of Radius R with Ω and we restrict the evolution to this domain. The requirements on the size of the ball will be specified later in this proof and the time T, up to when the evolution is considered, is such that the nodes of the network stay away from the boundary of Ω_R. Notice that if Ω is bounded domain and R is large enough the same definition can be made without altering Ω. Hence we will always refer to the domain as Ω_R.

Since the topological classes of $\mathcal{N}_1$ and $\mathcal{N}_2$ agree, Proposition 2.1 implies that the same coloring can be associated to both of them. From Theorem 3.3 that there are solutions u^1_ϵ and u^2_ϵ to (1)-(2)-(3) which as $\epsilon \to 0$ develop nodal sets that respectively agree with $\mathcal{N}_1$ and $\mathcal{N}_2$ in Ω_R and respect the associated coloring. Let ϕ^1_ϵ and ϕ^2_ϵ be the boundary conditions of u^1_ϵ and u^2_ϵ respectively. To simplify the notation, we assume that these functions have been already extended to the whole domain via a cut-off function. We denote

$$\tilde{\phi}^i_\epsilon(x,t) = \begin{cases} \chi_R(x)\phi^i_\epsilon(x,t) & \text{if } \Omega \text{ is bounded} \\ \phi^i_\epsilon(x,t) & \text{otherwise} \end{cases}.$$

Furthermore Lemma 3.1 implies (by choosing $w_\epsilon(x,t) = u^2_\epsilon(x,t) - \tilde{\phi}^1_\epsilon(x,t)$) that either

$$\sup_{\Omega_R \times [0,T_n]} |u^1_\epsilon - u^2_\epsilon| \to 0$$

or

$$\sup_{\Omega_R \times [0,T_n]} |u^1_\epsilon - u^2_\epsilon| \leq C \sup_{\Omega_R \times [0,T_n]} |F_\epsilon(u^2_\epsilon - \tilde{\phi}^1) - u^2_\epsilon + \tilde{\phi}^1|,$$

where

$$\begin{aligned} F_\epsilon(h) = &- \int_0^t \int_{\Omega_R} \mathcal{H}_{\Omega_R}(x,y,t-s) \left(\frac{\nabla_u W(h + \tilde{\phi}^1_\epsilon)}{\epsilon^2} + P(\tilde{\phi}^1_\epsilon) \right) (y,s) dyds \\ &+ \int_{\Omega_R} \mathcal{H}_{\Omega_R}(x,y,t)(\psi_\epsilon(y) - \tilde{\phi}^1_\epsilon(y,0))dy, \end{aligned}$$

Let us assume that we are in the second case. Since u_2 is a solution to (1) and $u^2_\epsilon(x,t) - \tilde{\phi}^2_\epsilon(x,t) = 0$ for $x \in \partial\Omega_R$ we have that

$$\begin{aligned} u^2_\epsilon(x,t) - \tilde{\phi}^2_\epsilon(x,t) = &- \int_0^t \int_{\Omega_R} \mathcal{H}_{\Omega}(x,y,t-s) \left(\frac{\nabla_u W(u^2_\epsilon)}{\epsilon^2} + P(\tilde{\phi}^2_\epsilon) \right) (y,s) dyds \\ &+ \int_{\Omega_R} \mathcal{H}_{\Omega}(x,y,t)(\psi_\epsilon(y) - \tilde{\phi}^2_\epsilon(y,0))dy, \end{aligned}$$

and

Notice that the functions ψ_ϵ coincide since the initial condition of the networks agree. Hence, we have:

$$\begin{aligned} (F_\epsilon(u^2_\epsilon - \tilde{\phi}^1_\epsilon) - u^2_\epsilon + \tilde{\phi}^1_\epsilon)(x,t) = &- \int_0^t \int_{\Omega_R} \mathcal{H}_{\Omega}(x,y,t-s) \left(P(\tilde{\phi}^1_\epsilon - \tilde{\phi}^2_\epsilon) \right) (y,s) dyds \\ &+ \int_{\Omega_R} \mathcal{H}_{\Omega}(x,y,t)(\tilde{\phi}^1_\epsilon(y,0) - \tilde{\phi i}^2_\epsilon(y,0))dy. \end{aligned}$$

Since, as $R \to \infty$ we have that $|\tilde{\phi}^1_\epsilon - \tilde{\phi}^2_\epsilon|_{C^{2,1}} \to 0$, we have that for any $\delta > 0$ there is an R such that

$$\sup_{(x,t) \in \Omega_R \times [0,T]} \left| F_\epsilon(u^2_\epsilon - \tilde{\phi}^2_\epsilon) - u^2_\epsilon - \tilde{\phi}^2_\epsilon \right| < \delta. \tag{39}$$

Now, if the evolutions of $\mathcal{N}_1$ and $\mathcal{N}_2$ are different we would have that $\sup_{\Omega \times [0,T_n]} |u^1_\epsilon - u^2_\epsilon| > \min_{i \neq j}\{|c_i - c_j|\}$. Choose δ such that $C\delta < \min_{i \neq j}\{|c_i - c_j|\}$ (where C is the constant given by Lemma 3.1). Then Lemma 3.1 and (39) imply:

$$\min_{i \neq j}\{|c_i - c_j|\} < \sup_{\Omega \times [0,T_n]} |u^1_\epsilon - u^2_\epsilon| \leq C \sup_{\Omega \times [0,T_n]} \leq \sup_{\Omega \times [0,T_n]} |F_\epsilon(u^2_\epsilon - \tilde{\phi}^2_\epsilon) - u^2_\epsilon + \tilde{\phi}^2_\epsilon| \leq C\delta,$$

which yields a contradiction. □

Remark 4.1. *We would like to remark that in the previous proof we assumed that the constant C in Lemma 3.1 can be chosen uniformly for every domain Ω_R. This in fact holds, since the bounds for $\tilde{\phi}_\epsilon$ and for u_ϵ provided by Lemma 2.2 in* [S2] *are uniform in $\mathbb{R}^2$.*

REFERENCES

[Br] **K. Brakke** *The motion of a surface by its mean curvature* Mathematical Notes # 20, Princeton University Press, Princeton (1978).

[BR] **Bronsard, L. and Reitich, F.** *On three-phase boundary motion and the singular limit of a vector-valued Ginzburg-Landau equation, Arch. Rational Mech. Anal. (124)*, No. 4 (1993), 355–379.

[MNT] **Carlo Mantegazza, Matteo Novaga, and Vincenzo Maria Tortorelli.** *Motion by curvature of planar networks. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **3** No. 2 (2004), 235–324.

[MS] **R.Mazzeo and M.Sáez.** *Self-similar expanding solutions for the planar network flow.* Seminaires et Congreses, Societe Mathematique de France.

[S1] **M. Sáez Trumper.** *Relaxation of the Flow of Triods via the vector-valued Parabolic Ginzburg-Landau equation.* To appear in J. Reine Angew. Math. (Crelle's Journal)

[S2] **M. Sáez-Trumper.** *Existence of a solution to a vector-valued Ginzburg-Landau equation with a three well potential. arXiv:math/0702662.*

[SS] **O. Schnürer and F. Schulze.** *Self-similar expanding networks to curve shortening flow.* arXiv:math-dg/0702698. To appear in Ann. Sc. Norm. Super. Pisa Cl. Sci.

[Sch] **O. Schnürer *et. al.*** *Evolution of convex shaped netwoks under curve shotening flow.* To appear in Trans. of the AMS.

MARIEL SÁEZ TRUMPER
EDIFICIO ROLANDO CHUAQUI. FACULTAD DE MATEMÁTICAS
PONTIFICIA UNIVERSIDAD CATÓLICA DE CHILE
AVDA. VICUÑA MACKENNA 4860, MACUL, SANTIAGO,
CHILE .
E-mail address: `mariel@mat.puc.cl`